



Australian Government

Bureau of Meteorology

BoM Efforts to Improve SSESs for AVHRR SST Level 3 Products

Christopher Griffin^{1,2}, Helen Beggs², Leon Majewski¹,

¹Observations & Engineering Branch, Bureau of Meteorology,
Australia

²CAWCR, Bureau of Meteorology, Australia

ST-VAL Breakout Session, 14th GHRSSST Science Team Meeting, Woods Hole, 17-21
June 2013



Point of Use

I'm interested in a typical or representative value

\bar{x}

I'm interested in a typical or representative bias

- Something that I can safely subtract from the value and get a better result in the long run.
- The supplier should be able to tell me if this will work.

μ

I'm interested in a standard deviation that tells how much uncertainty there might be in the reported value

- Maybe $z = \sqrt{n} \cdot \mu / \sigma$ would give me some useful information ?

σ

If the number of measurements that went into the determination is small, I'm interested to know that number


n





The Processing Chain

Match





Australian Government

Bureau of Meteorology

SSES L2P

sses_count

Number of *in situ* matches that contributed to the statistics

sses_bias

Estimate of the mean of the difference between the *in situ* and satellite measurement of SST

sses_standard_deviation


Estimate of the standard deviation of the difference between the *in situ* and satellite measurement of SST



SSES L2P

sses_count

- Still match based on common satellite zenith angle, quality level and day/night
- Use 1 year historical record as the baseline (when n is small)






Australian Government

Bureau of Meteorology

Processing L2P to L3U: Grid

- Grid

- Choose the *best* quality pixels for the overlap
- Weight the values, bias, and variance by the likelihood of providing a good estimate
- The likelihood is proportional to the size of the overlapping region
- Bias and variance are likelihood weighted
- A representative number of degrees of freedom is determined by scaling the likelihood




$$n_U = \frac{\sum_i w_i n_i}{\max_i w_i n_i}$$



SSES L3U

sses_count



ssses_count

“effective” number of L2P SSTs that
contributed to the statistics



Australian Government

Bureau of Meteorology

Processing L3U to L3C: Merge

- Merge – combining data from the same sensor
 - *best* quality pixels give a “best typical value” over the time period
 - Weight the values by the likelihood of providing a good estimate
 - The best typical value has the most supporting measurements and the lowest uncertainty (variance)
 - Number of degrees of freedom is scaled by the maximum likelihood (as before)
 - Equally weighted SST allow the standard deviation of the SST over time to be estimated



Australian Government
Bureau of Meteorology


SSES L3C

sses_count and sst_count

1 June 2013, NOAA-19, L3C 3 day (night) “best typical value” contains fewer degrees of freedom. Some measurements were included with lower contribution because of higher variance


sses_count

SSSES count
sses_count (count)



sst_count (count)

unweighted count of number of contributory SST measurements
sst_count (count)






SSES L3C

sea_surface_temperature and sst_mean

1 June 2013, NOAA-19, “best typical value” (left) compared to unweighted mean (right) over the 3 day (night) period





Australian Government

Bureau of Meteorology

Processing L3C to L3S: Merge

- Merge – combining data from the same sensor
 - When combining multiple L3C, consider all sources of measurement weighted by the count
 - Biases are subtracted before measurements are combined
 - The combined count is recorded



SSES L3S

sses_standard_deviation and sst_standard_deviation

1 June 2013, NOAA-19, L3S 6 day (night)

sses_standard_deviation


error relative to *in situ* measurements, of the “best typical” measurement

sst_standard_deviation

satellite measured variability


sses_standard_deviation (kelvin)

SSES standard deviation estimate
sses_standard_deviation (kelvin)



sst_standard_deviation (kelvin)

standard deviation estimate of contributory SST measurements
sst_standard_deviation (kelvin)





Australian Government

Bureau of Meteorology

Processing L3U to L3C/S: “Typical”

- “Typical”
 - Standard error of “best typical” SST scales with $1/\sqrt{n}$
 - *In situ* match up error contributes as before, not decreasing with the number of measurements
 - Keeping equally weighted SST statistics allows these components to be properly accounted for



Australian Government

Bureau of Meteorology

SSES L3C/S for multiple days

sst_count Number of measurements that contributed to the “best typical” value

sst_mean Equally weighted mean SST that contributed to the “best typical” value

sst_standard_deviation
Equally weighted standard deviation that contributed to the “best typical” value

Representative of detected diurnal variability if the files contain day and night SST



Australian Government

Bureau of Meteorology

What you should be able to do

- Use the `sea_surface_temperature` as a “best typical” value over the time scale implied by the data
- Use the bias to correct for deviations from *in situ* measurements
- Use $z = \sqrt{n} \cdot \mu / \sigma$ to give an idea about if the measurement is “unusual” or not
- Use n to inform the accuracy of μ and σ
- Use the unweighted statistics and the “best typical” value to quantify the variation seen over an extended time period



Australian Government

Bureau of Meteorology

... and what might be next !

- L2P SSES estimates could be more realistic
 - Satellite and Sun zenith angles, Latitude, Longitude...
- and L2P to L3U merging could build in the correlation between neighbouring SSES
- validation of the approach and the usefulness of bias and standard deviation as a correction to the SST should be done routinely
- we can produce long period L3S files which give indications of diurnal variation as well as typical values



Australian Government

Bureau of Meteorology

Thank you...

Christopher Griffin

C.Griffin@bom.gov.au



Australian Government

Bureau of Meteorology

What this is about

- Rationalizing the current approach to make a little more sense from a *statistical* and *point of use* point of view.
- Respecting the *differences* in the way that data is combined in the formation of GHR SST SSES.
- Providing a *essential* set of fields in GHR SST files that may help to understand the uncertainties of measurement.
- Giving information which should be closer to being really usable.



Australian Government

Bureau of Meteorology

What this is not about

- A redefinition of SSES within GHRSSST SST products.

We are more interested in better communicating and encouraging the “point(s) of use”.
- A fundamental change to the way L2P SSES are currently estimated within ABOM products.

We have not considered changes to this apart from extending historical horizons to increase the number of *in situ* matches.
- A major change or “best” statistical solution to the problem.



Australian Government

Bureau of Meteorology

Processing L2: Match

- Match
 - Use *in situ* buoy measurements to match SST measurements, look at the differences
 - Bin the differences in critical parameters (satellite zenith angle, quality level, day / night)
 - If we don't have enough data, reference historical information (past year) to make realistic estimates
 - Report the number of measurements used in the estimate