

# Application and Evaluation of Diurnal Warming Models Forced with GFS Model Inputs

Gary A. Wick  
NOAA ESRL/PSD

# Motivation

- Preparation for real-time computation of diurnal warming estimates
- Facilitate further comparison and validation of models

# Components

- Model Inputs
  - GFS analysis fields, 6 hourly, 0.5 degree
    - Wind stress
    - Radiative and turbulent fluxes
  - Wave Watch III Wave Model
    - Wave period, direction, and significant wave height
- Models currently implemented
  - Kantha Clayson
  - COARE
  - Parameterizations


# Logistics

- Models initialized based on SST
- Fluxes interpolated to model time step
- Model run globally for 2 days with output taken from the second day
- Real-time implementation just being finalized
- Warming compared against SEVIRI observations

# **SAMPLE MODEL RESULTS**


# Kantha Clayson Stokes Drift Model

## Diurnal Warming Evolution, 00-23 UTC, 21 March 2013



# Gentemann Parameterization

## Diurnal Warming Evolution, 00-23 UTC, 21 March 2013




# **COMPARISON OF WARMING DISTRIBUTIONS WITH SEVIRI OBSERVATIONS**


# Comparison of Observed and Modeled Diurnal Warming

## SEVIRI Domain – 21 March 2013, 1400 UTC


### K-C Stokes Drift




### K-C Stokes Enh (2)




### K-C Stokes Enhanced




### Wick K-C Blended



### COARE




### Gentemann




# Comparison of Observed and Modeled Diurnal Warming

## SEVIRI Domain – 21 March 2013, 2200 UTC


### K-C Stokes Drift




### K-C Stokes Enh (2)




### K-C Stokes Enhanced




### Wick K-C Blended



### COARE



### Gentemann




# **TIME EVOLUTION OF WARMING DISTRIBUTIONS FOR MODELS**

# Comparison of Observed and Modeled Diurnal Warming


## SEVIRI Domain – 21 March 2013

### Kantha Clayson Model with Stokes Drift


**0200 UTC**




**0600 UTC**




**1000 UTC**




**1400 UTC**



**1800 UTC**



**2200 UTC**




# Summary

- Automated daily computation of global diurnal warming from NWP forcing coming on line
- Display, distribution, and evaluation of products to follow


# COARE Model

Diurnal Warming Evolution, 00-23 UTC, 21 March 2013




# Kantha Clayson Stokes (Enhanced v2) Model

Diurnal Warming Evolution, 00-23 UTC, 21 March 2013



# Wick Blended Kantha Clayson Model

## Diurnal Warming Evolution, 00-23 UTC, 21 March 2013




# Comparison of Observed and Modeled Diurnal Warming


SEVIRI Domain – 21 March 2013

Kantha Clayson Model with Stokes Drift and Modified Constants (v2)


0200 UTC




0600 UTC




1000 UTC




1400 UTC



1800 UTC



2200 UTC




# Comparison of Observed and Modeled Diurnal Warming


## SEVIRI Domain – 21 March 2013

### Wick Blended Kantha Clayson Model


**0200 UTC**




**0600 UTC**




**1000 UTC**




**1400 UTC**



**1800 UTC**



**2200 UTC**




# Comparison of Observed and Modeled Diurnal Warming


## SEVIRI Domain – 21 March 2013

### COARE Model


**0200 UTC**




**0600 UTC**




**1000 UTC**




**1400 UTC**



**1800 UTC**



**2200 UTC**




# Comparison of Observed and Modeled Diurnal Warming


## SEVIRI Domain – 22 March 2013

### Kantha Clayson Model with Stokes Drift


**0200 UTC**




**0600 UTC**




**1000 UTC**



**1400 UTC**



**1800 UTC**



**2200 UTC**

